ANAPHYLAXIS EMERGENCY ACTION PLAN

NAME:	AGE:	_
ALLERGY TO:		
Asthma		
Concurrent medications, if any:		
SYMPTOMS OF ANAPHYLAXIS INCLUDE:		
	swelling of lips and/or tongue	
<u>-</u>	3, 19, 11, 11, 11, 11, 11, 11, 11, 11, 11	
GUT vomiting, diarrhea, cramps		
	ss of breath, cough, wheeze llse, dizziness, passing out	
Only a few symptoms may be present. Severity of symptoms can change quickly.		
*Some symptoms can be life-threatening! ACT FAST!		
WHAT TO DO:	ank ana).	☐ Turinia et 0.45 mm
1. INJECT EPINEPHRINE IN THIGH USING (ch	EpiPen (0.3 mg)	Twinject 0.15 mg
Other medication/dose/route:		
IMPORTANT: ASTHMA PUFFERS AND/OR ANTIHISTAMINES CAN'T BE DEPENDED ON IN ANAPHYLAXIS!		
2. CALL 911 or RESCUE SQUAD (BEFORE CA	LLING CONTACTS)!	
3. Emergency contact #1: home	work	cell
Emergency contact #2: home	work	cell
Emergency contact #3: home	work	cell
DO NOT HESITATE TO GIVE EPINEPHRINE!		
COMMENTS:		
Doctor's Signature/Date Parent's Signature (for individuals under age 18 yrs)/Date		